

# Cabral®

24 Coated Tablets

## Formula:

1 Coated tablet

Active Substance : 400 mg Phenyramidol HCl

Excipient : Tartazine, titanium dioxide, grounded sugar.

## Pharmacological Properties:

Cabral breaks pain-strain chain by blocking polysynaptic reflexes in brain and medulla spinalis. In this manner, it relieves muscle tension, and it has a strong analgesic effect in muscle pain. It does not affect monosynaptic reflexes.

As a myorelaxant and analgesic, Cabral is indicated in the treatment of acute and chronic pains in striated muscle and the other members of movement system.

## Indications:

Waist pains, lumbago, sciatica, disk disorders and supportive treatment to other medico-mechanic or chiropractic therapies such as under water massages. Relieves muscle tension and pain, and maintains the function of cramped skeletal muscle.

## Contraindications:

It should not be administered in individuals who are sensitive to any substance of the preparation composition.

## Special Warnings and Precautions for Use:

It is recommended to discontinue Cabral and consult to a doctor in cases including hemorrhage when concomitant administration with anticoagulants, alteration in carbohydrate metabolism with antidiabetics, eye tremor with anticonvulsives, vision disorders, appearance of uncoordinated movements.

Although the amount of drug colorant Tartazine is lesser than the limits, it is recommended to use cautiously because of the hypersensitivity reactions.

## Side Effects/Advers Effects:

It may cause bloated feeling in the stomach or nausea, burn feeling in tongue or skin exanthema in some hypersensitive patients. However these are temporary, also numbness, pruritus and rash may be occurred. Drug should be discontinued when hypersensitivity and rash occur.

## CONSULT YOUR DOCTOR IF AN UNEXPECTED EFFECT ENSUES

## Drug Interactions and Other Forms of Interaction:

Cabral coated tablets must be administered with caution in the patients who are treated with cumarin derivatives (anticoagulants), tolbutamide, oral antidiabetics and diphenylhydantoin.

## Administration Route and Dosage:

Three times a day 1-2 coated tablets are swallowed without chewing after meals.

## Overdose and Treatment:

Daily doses up to 3200 mg are well tolerated. However, it is recommended to consult to doctor in case of any symptoms related to the drug.

## Type of Supply and Container Content:

Blisters containing 24 coated tablets.

## Storage Conditions:

Store at room temperatures less than 25 °C.

## Other Pharmaceutical Dosage Forms Present in the Market:

3 ml x 3 ampoules.

Keep this and all drugs out of reach of children and store in its packaging.  
Do not use without consulting your doctor.

**Registration Holder: Dr. F. Frik İlaç San. Ve Tic. A.Ş. Küçükyalı / İSTANBUL**  
**Place of Manufacture: Yeni İlaç ve Hammaddeleri San. Tic. A.Ş. Esenyurt / İSTANBUL**  
**Registration Date and No.: 11/11/1970 – 106/12**  
Sold with prescription only.